

# Research Statement

Ali Alilooee

My primary field of research is commutative algebra. I am especially interested in problems in commutative algebra which can be translated into the language of combinatorics or algebraic geometry. One of the most useful techniques applied to connect commutative algebra to combinatorics is assigning a squarefree monomial ideal to a graph or a simplicial complex to make a dictionary between their algebraic and combinatorial properties.

A squarefree monomial is a product of distinct variables, for example  $x_1x_3x_4$  in the polynomial ring  $k[x_1, \dots, x_4]$ . A squarefree monomial ideal is an ideal generated by squarefree monomials. The central theme of my research is the study of squarefree monomial ideals from a combinatorial perspective.

Here are some specific areas which I am currently studying. A more detailed description for each follows after.

- **Rees algebra of squarefree monomial ideals:**

The equations of Rees algebra of quadratic squarefree monomial ideals are known. The focus of my work is to find a combinatorial interpretation for equations of Rees algebras of squarefree monomial ideals of higher degrees.

- **Regularity of powers of edge ideals:**

In [2] with collaboration with A. Banerjee we investigated the asymptotic behavior of regularity of a class of bipartite graphs. Banerjee, Beyarslan, Hà and I plan to do more with improving the asymptotic formula for the edge ideals of trees and cycles in [16] to a more general case.

- **$j$ -multiplicity of edge ideals:**

$j$ -multiplicity is a generalization of Hilbert-Samuel multiplicity for ideals which are not primary. I am working on  $j$ -multiplicity of edge ideals of a graph. This is joint work with J. Validashti.

Below I describe the principal themes and techniques for each of these projects.

# 1 Rees Algebra of Squarefree Monomial Ideals

Let  $I = (f_1, \dots, f_q)$  be a monomial ideal in a polynomial ring  $R = k[x_1, \dots, x_n]$  over a field  $k$ . We denote the **Rees algebra** of  $I$  by  $R[It] = R[f_1t, \dots, f_qt]$  and consider the homomorphism  $\psi$  of algebras

$$\psi : R[T_1, \dots, T_q] \longrightarrow R[It], \quad T_i \mapsto f_it$$

where  $T_i$  are indeterminate. If  $J$  is kernel of  $\psi$ , then  $R[It] = S/J$  where  $S = R[T_1, \dots, T_q]$ . The ideal  $J$  is called **defining ideal** of  $R[It]$  and its minimal generators are called the **Rees equations** of  $I$ . The defining ideal  $J$  of  $R[It]$  is graded. In other words, we have

$$J = J_1 \oplus J_2 \oplus \dots$$

where  $J_s$  for  $s \geq 1$  is  $R$ -modules. These equations carry a lot of information about  $R[It]$ ; see for example [17], for more details.

An ideal  $I$  is said to be **of linear type** if  $J = (J_1)$ ; in other words, the defining ideal of  $R[It]$  is generated by linear combinations of variables  $T_1, \dots, T_q$ .

Rees algebras and ideals of linear type have been investigated by many authors (see for example [4], [5], [9], [12],[13])

We can define  $T_{\alpha,\beta}(I) \in J_s$  as a **redundant Rees equation** of  $I$ , if we have

$$T_{\alpha,\beta}(I) \in J_1S + \dots + J_{s-1}S.$$

To identify ideals of linear type, it is enough to characterize redundant equations.

I am interested in giving a combinatorial interpretation of irredundant  $T_{\alpha,\beta}(I)$  for a squarefree monomial ideal  $I$ . The simplest case of a squarefree monomial ideal is an edge ideal.

In 1995 Villarreal gave a combinatorial characterization of irredundant generators for edge ideals of graphs [18] by attributing irredundant generators of  $J_s$  to closed even walks. Motivated by this work I defined **simplicial closed even walks** in my thesis for simplicial complexes to generalize graph case.

Faridi and I proved in [3] that if  $T_{\alpha,\beta}(I)$  is an irredundant generator of  $J_s$ , then the generators of  $I$  involved in  $T_{\alpha,\beta}(I)$  form a simplicial even walk. Our class of simplicial even walks includes *even special cycles* (see [10]) as known in hypergraph theory.

By using the concept of the simplicial closed even walks, we can give a necessary condition for a squarefree monomial ideals of linear type [3].

**Theorem 1.1.** *Let  $I$  be a squarefree monomial ideal and suppose the facet complex  $\mathcal{F}(I)$  has no simplicial closed even walk. Then  $I$  is of linear type.*

Also in my thesis I showed that every simplicial closed even walk contains a simplicial cycle (not necessarily of the same length).

By using this new result we can conclude every simplicial tree is of linear type. (Simplicial trees are generalization of trees in graph theory. For more details see [7].) This fact can also be deduced by the concept of  $M$ -sequences [4].

## 1.1 Further Work

Generally the converse of Theorem 1.1 is not correct. My goal is to further investigate the structure of simplicial closed even walk so that I can give a more effective criterion to replace Theorem 1.1.

**Question 1.** Find a more effective criterion for squarefree monomial ideals of linear type.

Fouli and Lin [9] used the line graphs of squarefree monomial ideals to give a simple test for ideals of linear type. In my thesis I improved their test as demonstrated in the following theorem.

**Theorem 1.2.** *Let  $I$  be a squarefree monomial ideal and  $\Delta$  be its facet complex. If  $L(\Delta)$  (line graph of  $\Delta$ ) contains no even cycle, then  $I$  is of linear type.*

One can ask what happens for line graphs which contain an even cycle. In general we can say nothing about linearity of  $I$  by using these line graphs. I am currently working on improving Theorem 1.2.

## 2 Regularity of Powers of Edge Ideals

The **Castelnuovo-Mumford regularity** of  $R/I$ , written as  $\text{reg}(R/I)$ , is given by

$$\text{reg}(R/I) := \max\{j - i \mid \beta_{i,j}(R/I) \neq 0\}.$$

Asymptotic behavior of the regularity is one of the interesting ongoing research areas in commutative algebra. The following theorem by Cutkosky, Herzog and Trung is very motivating for many researchers.

**Theorem 2.1** (Theorem 1.1, [6]). *Let  $I$  be a homogeneous ideal, the regularity  $I^n$  is a linear function for  $n \gg 0$ .*

Determining the exact linear function of  $\text{reg}(I^n)$  and evaluating  $n$  for which  $\text{reg}(I^n)$  becomes linear, have been studied by many researchers.

In my research I consider this question for edge ideal of a simple graph. More precisely if  $G$  is a simple graph whose edge ideal is denoted by  $I(G)$ , I am interested in finding values  $b$  and  $n_0$  for which we have

$$\text{reg}(I(G)^n) = 2n + b \quad \text{for } n \geq n_0.$$

The values  $b$  and  $n_0$  have been computed for a few classes of graphs. These classes are as follows

- **Herzog, Hibi and Zheng [11]:** If  $\text{reg}(I(G)) = 2$  ( $I(G)$  has linear resolution), then  $\text{reg}(I(G)^n) = 2n$  for all  $n \geq 1$  ( $b = 0, n_0 = 1$ ).
- **Alilooee and Banerjee [2]:** If  $G$  is a bipartite graph and  $\text{reg}(I(G)) = 3$ , then  $\text{reg}(I(G)^n) = 2n + 1$  for all  $n \geq 1$  ( $b = 1, n_0 = 1$ ).
- **Beyarslan, Hà and Trung [16]:** If  $G$  is a forest, then  $\text{reg}(I(G)^n) = 2n + \nu(G) - 1$  for all  $n \geq 1$  ( $b = \nu(G) - 1, n_0 = 1$ ) where  $\nu(G)$  denotes the maximum induced matching number of  $G$ .
- **Beyarslan, Hà and Trung [16]:**  $\text{reg}(I(C_m)^n) = 2n + \nu(C_m) - 1$  for all  $n \geq 2$  ( $b = \nu(C_m) - 1, n_0 = 2$ ).
- **Ferrò, Murgia and Olteanu [8]** If  $I(G)$  is an initial and final lexsegment edge ideal, then  $\text{reg}(I(G)^n) = 2n$  for all  $n \geq 1$  ( $b = 0, n_0 = 1$ ).

Beyarslan, Hà and Trung in [16] have shown that if  $G$  is a graph with edge ideal  $I(G)$  and  $\nu(G)$  denotes the induced matching number of  $G$ , then we have

$$\text{reg}(I(G)^n) \geq 2n + \nu(G) - 1 \quad \text{for } n \geq 1.$$

They asked the following question.

**Question 2.** If  $G$  is a simple graph whose edge ideal is  $I = I(G)$ , then is it true that  $\text{reg}(I(G)^n) = 2n + \nu(G) - 1$  for all  $n \geq \text{reg}(I(G)) - 1$ ?

Beyarslan, Hà and Trung showed that this question is true for forests and cycles. More recently, Banerjee, Beyarslan, Hà and I showed that in general Question 2 is not true.

We intend to identify the classes of graphs for which Question 2 is true. It suffices to identify classes with  $\text{reg}(I(G)^n) \leq 2n + \nu(G) - 1$  for  $n \gg 0$ . Banerjee, Beyarslan, Hà and I have shown that for a graph  $G$  we have

$$\text{reg}(I(G)^n) \leq 2n + \text{reg}(I(G)) - 2 \quad \text{for } n \geq 1. \tag{2.1}$$

By using this upper bound we can conclude that Question 2 holds for the classes of graphs such that  $\text{reg}(I(G)) = \nu(G) + 1$ . For the following classes of graphs, the regularity of the edge ideal is known to be  $\nu(G) + 1$  and so regularity of powers can be computed.

- **Chordal graphs** (Hà, Van Tuyl (2008)).
- **Sequentially Cohen-Macaulay bipartite graphs** (Van Tuy (2009)).

- **Unmixed bipartite graph** (Kummini (2009)).
- **Very well-covered graphs** (Mahmoudi, Mousivand, Crupi, Rinaldo, Terai, Yassemi (2011)).
- **$(C_4, C_5)$ -free vertex decomposable graphs** (Biyikoglu, Civan (2012)).
- **$C_5$ -free vertex decomposable graphs** (Khosh-Ahang, Moradi (2013)).
- **Unicyclic graphs** (Alilooee, Beyarslan (2015)).

## 2.1 Further Work

I would like to add more classes of graphs to the above list. Currently, I am working on the graphs with two cycles. I am also interested in bipartite graphs. For these graphs Há and I have made the following conjecture.

**Conjecture 1.** *Let  $G$  be a bipartite graph with edge ideal  $I(G)$  and induced matching number  $\nu(G)$ . Then*

$$\nu(G) + 1 \leq \text{reg}(I(G)) \leq \nu(G) + 2.$$

I am also working on a similar version of equation 2.1 for hypergraphs. In other words I would like to see that if the equation 2.1 is true for each squarefree monomial ideal.

## 3 $j$ -Multiplicity of Edge Ideals

Hilbert-Samuel multiplicity is one of the most important invariants in homological algebra, commutative algebra and algebraic geometry. This notation was defined in 1954 by D. Northcott and D. Rees [15]. The Hilbert-Samuel multiplicity is defined in a local ring for  $\mathfrak{m}$ -primary ideals where  $\mathfrak{m}$  is the maximal ideal. In 1993  $j$ -multiplicity was defined by R. Achilles and M. Manaresi [1] to extend Hilbert-Samuel multiplicity notation for an ideal that is not necessarily  $\mathfrak{m}$ -primary.

We are ready to state the definition of  $j$ -multiplicity of an ideal.

**Definition 3.1.** Let  $(R, \mathfrak{m}, k)$  be a Noetherian local ring of dimension  $d$  and  $I$  be an ideal in  $R$ . We call the positive integer given by

$$j(I) = \lim_{n \rightarrow \infty} \frac{(d-1)! \lambda(H_{\mathfrak{m}}^0(I^n/I^{n+1}))}{n^{d-1}}.$$

the  $j$ -multiplicity of  $I$  in  $R$ .

We have the following properties for  $j$ -multiplicity (see [1])

- $j(I) \neq 0 \iff \ell(I) = \dim(R)$  where  $\ell(I)$  denoted the analytic spread of  $I$  (The Krull dimension of the fiber cone of  $I$ ).
- If  $I$  is an  $m$ -primary ideal in  $R$ , then we have

$$j(I) = e(I, R) = e(\text{Gr}_I(R))$$

where  $e(-)$  denotes the Hilbert-Samuel multiplicity.

- For a monomial ideal  $I$  in the polynomial ring  $R = k[x_1, \dots, x_d]$ , we have

$$j(I) = d! \text{vol}(\text{pyr}(I)).$$

where  $\text{pyr}(I)$  is a union of pyramid of bounded faces of Newton polyhedron of  $I$ . (see Theorem 3.2 in [14])

In general computing  $j$ -multiplicity of an ideal is not easy. In my work with J. Validashti, we have studied  $j$ -multiplicity of edge ideal of a graph. Our results are described below.

### 3.1 Edge Ideals

Let  $G$  be a simple graph and  $P(G)$  be the Newton Polyhedron of the edge ideal  $I(G)$ . To compute  $j$ -multiplicity of the edge ideal of  $G$  from Theorem 3.2 in [14] we need to consider bounded faces of  $P(G)$ . The convex hull of the vertices of  $P(G)$ , called the **edge Polytope**  $F(G)$  of  $G$ , is the only bounded face.

By using this fact I in collaboration with Validashti proved the following theorem to classify graphs with non-zero  $j$ -multiplicity.

**Remark 3.2.**  $j(I(G)) \neq 0$  if and only if no connected component of  $G$  is bipartite.

To compute  $j$ -multiplicity we first consider connected unicycle graphs in which obviously the number of vertices and edges are the same. For a unicycle graph we have the following formula to compute  $j$ -multiplicity of its edge ideal. This formula has been computed by Jeffries and Montaña in [14] by using different method.

**Corollary 3.3.** *If  $G$  is a graph which its components are unicycle, then*

$$j(I(G)) = 2^c \alpha(G)$$

where  $c$  is the number of connected component of  $G$  and  $\alpha(G) = 1$  if  $G$  has no bipartite connected component, and zero otherwise .

By using  $K[G]$ , the edge subrings of a graph  $G$ , Validashi and I showed the following theorem.

**Theorem 3.4.** *If  $G$  is a simple graph with  $c(G)$  connected components, then*

$$j(I(G)) = 2^{c(G)}e(K[G]).$$

By using this theorem we have the following interesting formula for  $j$ -multiplicity of complete graphs.

**Corollary 3.5.** *If  $G$  is a complete graph with  $n$  vertices, we have*

$$j(I(G)) = 2(2^{n-1} - n) = 2^n - 2n.$$

**Question 3.** We gave formulas to compute  $j$ -multiplicity for some classes of graphs (trees, unicycles, bipartite and complete). Is there any other classes of graphs which their  $j$ -multiplicity can be given by a formula?

**Question 4.** What is the interpretation of  $2^n - 2n$  for complete graphs with  $n$  vertices?

**Question 5.** If  $\ell(I(G)) \neq \dim R$ , can we say  $e(K[G])$  is the next generalized Hilbert coefficient of  $I(G)$ ?

## References

- [1] R. Achilles and M. Manaresi. Multiplicity for ideals of maximal analytic spread and intersection theory. *J. Math. Kyoto Univ.*, 33(4):1029–1046, 1993.
- [2] A. Alilooee and A. Banerjee. Powers of edge ideals of regularity three bipartite graphs. *arXiv:1408.2557 [math.AC]*, 2014.
- [3] A. Alilooee and S. Faridi. When is a squarefree monomial ideal of linear type. *arXiv:1309.1771 [math.AC]*, 2013.
- [4] A. Conca and E. De Negri.  $M$ -sequences, graph ideals, and ladder ideals of linear type. *J. Algebra*, 211(2):599–624, 1999.
- [5] D. L. Costa. Sequences of linear type. *J. Algebra*, 94(1):256–263, 1985.
- [6] S.D. Cutkosky, J. Herzog, and N. V. Trung. Asymptotic behaviour of the Castelnuovo-Mumford regularity. *Compositio Math.*, 118(3):243–261, 1999.
- [7] S. Faridi. Simplicial trees are sequentially cohen-macaulay. *J. Pure Appl. Algebra*, 190(1-3):121–136, 2004.
- [8] Carmela Ferrò, Mariella Murgia, and Oana Olteanu. Powers of edge ideals. *Matematiche (Catania)*, 67(1):129–144, 2012.

- 
- [9] L. Fouli and K. N. Lin. Rees algebras of square-free monomial ideals. *arXiv:1205.3127v2 [math.AC]*, 2013.
- [10] J. Herzog, T. Hibi, N. V. Trung, and X. Zheng. Standard graded vertex cover algebras, cycles and leaves. *Trans. Amer. Math. Soc.*, 360(12):6231–6249, 2008.
- [11] J. Herzog, T. Hibi, and X. Zheng. Monomial ideals whose powers have a linear resolution. *Math. Scand.*, 95(1):23–32, 2004.
- [12] C. Huneke. The theory of  $d$ -sequences and powers of ideals. *Adv. in Math.*, 46(3):249–279, 1982.
- [13] C. Huneke. Determinantal ideals of linear type. *Arch. Math. (Basel)*, 47(4):324–329, 1986.
- [14] J. Jeffries and J. Montaña. The  $j$ -multiplicity of monomial ideals. *Math. Res. Lett.*, 20(4):729–744, 2013.
- [15] D. G. Northcott and D. Rees. Reductions of ideals in local rings. *Proc. Cambridge Philos. Soc.*, 50:145–158, 1954.
- [16] T. N. Trung S. Beyarslan, H. T. Ha. The regularity of powers of edge ideals. *arXiv:1409.0277 [math.AC]*, 2014.
- [17] W. Vasconcelos. *Arithmetic of Blowup Algebras*. Cambridge University Press, Cambridge, 1994.
- [18] R. H. Villarreal. Rees algebras of edge ideals. *Comm. Algebra*, 23(9):3513–3524, 1995.